



**University of  
Zurich**<sup>UZH</sup>

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2010

---

## **THE (1405), (1475), f1(1420), AND f1(1510) (Revised February 2010)**

Amsler, C ; Masoni, A

DOI: <https://doi.org/10.1088/0954-3899/37/7A/075021>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-46085>

Journal Article

Accepted Version

Originally published at:

Amsler, C; Masoni, A (2010). THE (1405), (1475), f1(1420), AND f1(1510) (Revised February 2010).

Journal of Physics G: Nuclear and Particle Physics, 37(7A):680-684.

DOI: <https://doi.org/10.1088/0954-3899/37/7A/075021>

## THE $\eta(1405)$ , $\eta(1475)$ , $f_1(1420)$ , AND $f_1(1510)$

Revised February 2010 by C. Amsler (Zürich) and A. Masoni (INFN Cagliari).

The first observation of the  $\eta(1440)$  was made in  $p\bar{p}$  annihilation at rest into  $\eta(1440)\pi^+\pi^-$ ,  $\eta(1440) \rightarrow K\bar{K}\pi$  [1]. This state was reported to decay through  $a_0(980)\pi$  and  $K^*(892)\bar{K}$  with roughly equal contributions. The  $\eta(1440)$  was also observed in radiative  $J/\psi(1S)$  decay into  $K\bar{K}\pi$  [2–4] and  $\gamma\rho$  [5]. There is now evidence for the existence of two pseudoscalars in this mass region, the  $\eta(1405)$  and  $\eta(1475)$ . The former decays mainly through  $a_0(980)\pi$  (or direct  $K\bar{K}\pi$ ) and the latter mainly to  $K^*(892)\bar{K}$ .

The simultaneous observation of two pseudoscalars is reported in three production mechanisms:  $\pi^-p$  [6,7]; radiative  $J/\psi(1S)$  decay [8,9]; and  $\bar{p}p$  annihilation at rest [11–14]. All of them give values for the masses, widths, and decay modes in reasonable agreement. However, Ref. [9] favors a state decaying into  $K^*(892)\bar{K}$  at a lower mass than the state decaying into  $a_0(980)\pi$ , although agreement with MARK-III is not excluded. In  $J/\psi(1S)$  radiative decay, the  $\eta(1405)$  decays into  $K\bar{K}\pi$  through  $a_0(980)\pi$ , and hence a signal is also expected in the  $\eta\pi\pi$  mass spectrum. This was indeed observed by MARK III in  $\eta\pi^+\pi^-$  [15], which reports a mass of 1400 MeV, in line with the existence of the  $\eta(1405)$  decaying into  $a_0(980)\pi$ . BES [10] reports an enhancement in  $K^+K^-\pi^0$  around 1.44 GeV in  $J/\psi(1S)$  decay, recoiling against an  $\omega$  (but not a  $\phi$ ) without resolving the presence of two states nor performing a spin-parity analysis, due to low statistics. This state could also be the  $f_1(1420)$  (see below).

The  $\eta(1405)$  is also observed in  $\bar{p}p$  annihilation at rest into  $\eta\pi^+\pi^-\pi^0\pi^0$ , where it decays into  $\eta\pi\pi$  [16]. The intermediate  $a_0(980)\pi$  accounts for roughly half of the  $\eta\pi\pi$  signal, in agreement with MARK III [15] and DM2 [4].

The  $\eta(1295)$  has been observed by four  $\pi^-p$  experiments [7,17–19], and evidence is reported in  $\bar{p}p$  annihilation [23–25]. In  $J/\psi(1S)$  radiative decay, an  $\eta(1295)$  signal is evident in the  $0^{-+}$   $\eta\pi\pi$  wave of the DM2 data [9]. Also BaBar [20] reports evidence for a signal around 1295 MeV in  $B$  decays into  $\eta\pi\pi K$ .

However, the existence of the  $\eta(1295)$  is questioned in Refs. [21] and [22]. The authors claim a single pseudoscalar meson in the 1400 MeV region. This conclusion is based on properties of the wave functions in the  $^3P_0$  model (and on an unpublished analysis of the annihilation  $\bar{p}p \rightarrow 4\pi\eta$ ). The pseudoscalar signal around 1400 MeV is then attributed to the first radial excitation of the  $\eta$ .

Assuming establishment of the  $\eta(1295)$ , the  $\eta(1475)$  could be the first radial excitation of the  $\eta'$ , with the  $\eta(1295)$  being the first radial excitation of the  $\eta$ . Ideal mixing, suggested by the  $\eta(1295)$  and  $\pi(1300)$  mass degeneracy, would then imply that the second isoscalar in the nonet is mainly  $s\bar{s}$ , and hence couples to  $K^*\bar{K}$ , in agreement with properties of the  $\eta(1475)$ . Also, its width matches the expected width for the radially excited  $s\bar{s}$  state [26,27]. A study of radial excitations of pseudoscalar mesons [28] favors the  $s\bar{s}$  interpretation of the  $\eta(1475)$ . However, due to the strong kinematical suppression the data are not sufficient to exclude a sizeable  $s\bar{s}$  admixture also in the  $\eta(1405)$ .

The  $K\bar{K}\pi$  and  $\eta\pi\pi$  channels were studied in  $\gamma\gamma$  collisions by L3 [29]. The analysis leads to a clear  $\eta(1475)$  signal in  $K\bar{K}\pi$ , decaying into  $K^*\bar{K}$ , very well identified in the untagged data sample, where contamination from spin 1 resonances is not allowed. At the same time, L3 [29] did not observe the  $\eta(1405)$ , neither in  $K\bar{K}\pi$  nor in  $\eta\pi\pi$ . The observation of the  $\eta(1475)$ , combined with the absence of an  $\eta(1405)$  signal, strengthens the two-resonances hypothesis. Since gluonium production is presumably suppressed in  $\gamma\gamma$  collisions, the L3 results [29] suggest that  $\eta(1405)$  has a large gluonic content (see also Refs. [30] and [31]).

The L3 result is somewhat in disagreement with that of CLEO-II, which did not observe any pseudoscalar signal in  $\gamma\gamma \rightarrow \eta(1475) \rightarrow K_S^0 K^\pm \pi^\mp$  [32]. However, more data are required. Moreover, after the CLEO-II result, L3 performed a further analysis with full statistics [33], confirming the evidence of the  $\eta(1475)$  observed by L3. The CLEO upper limit [32] for  $\Gamma_{\gamma\gamma}(\eta(1475))$ , and the L3 results [33], are consistent with the world average for the  $\eta(1475)$  width.

BaBar [20] also reports the  $\eta(1475)$  in  $B$  decays into  $K\bar{K}\pi$  (and possibly  $\eta\pi\pi$ ). Upper limits are given for  $\eta(1405)$  decay into  $K\bar{K}^*$ . The data sample is not sufficient to identify a possible  $\eta(1405)$  contribution into  $\eta\pi\pi$ .

The gluonium interpretation for the  $\eta(1405)$  is not favored by lattice gauge theories which predict the  $0^{-+}$  state above 2 GeV [34]. However, the  $\eta(1405)$  is an excellent candidate for the  $0^{-+}$  glueball in the fluxtube model [35]. In this model, the  $0^{++}$   $f_0(1500)$  glueball is also naturally related to a  $0^{-+}$  glueball with mass degeneracy broken in QCD. Also, Ref. [36] shows that the pseudoscalar glueball could lie at a lower mass than predicted from lattice calculation. In this model the  $\eta(1405)$  appears as the natural glueball candidate (see also Refs. [37] and [38]). A detailed review of the experimental situation is available in Ref. [39].

Let us now deal with  $1^{++}$  isoscalars. The  $f_1(1420)$ , decaying into  $K^*\bar{K}$ , was first reported in  $\pi^-p$  reactions at 4 GeV/c [40]. However, later analyses found that the 1400–1500 MeV region was far more complex [41–43]. A reanalysis of the MARK III data in radiative  $J/\psi(1S)$  decay into  $K\bar{K}\pi$  [8] shows the  $f_1(1420)$  decaying into  $K^*\bar{K}$ . Also, a  $C=+1$  state is observed in tagged  $\gamma\gamma$  collisions (*e.g.*, Ref. [44]).

In  $\pi^-p \rightarrow \eta\pi\pi n$  charge-exchange reactions at 8–9 GeV/c the  $\eta\pi\pi$  mass spectrum is dominated by the  $\eta(1440)$  and  $\eta(1295)$  [17,45], and at 100 GeV/c Ref. [18] reports the  $\eta(1295)$  and  $\eta(1440)$  decaying into  $\eta\pi^0\pi^0$  with a weak  $f_1(1285)$  signal, and no evidence for the  $f_1(1420)$ .

Axial ( $1^{++}$ ) mesons are not observed in  $\bar{p}p$  annihilation at rest in liquid hydrogen, which proceeds dominantly through  $S$ -wave annihilation. However, in gaseous hydrogen,  $P$ -wave annihilation is enhanced and, indeed, Ref. [12] reports  $f_1(1420)$  decaying into  $K^*\bar{K}$ . The  $f_1(1420)$ , decaying into  $K\bar{K}\pi$ , is also seen in  $pp$  central production, together with the  $f_1(1285)$ . The latter decays via  $a_0(980)\pi$ , and the former only via  $K^*\bar{K}$ , while the  $\eta(1440)$  is absent [46,47]. The  $K_S K_S \pi^0$  decay mode of the  $f_1(1420)$  establishes unambiguously  $C=+1$ . On the other hand, there is no evidence for any state decaying into  $\eta\pi\pi$  around

1400 MeV, and hence the  $\eta\pi\pi$  mode of the  $f_1(1420)$  must be suppressed [48].

We now turn to the experimental evidence for the  $f_1(1510)$ . Two states, the  $f_1(1420)$  and  $f_1(1510)$ , decaying into  $K^*\bar{K}$ , compete for the  $s\bar{s}$  assignment in the  $1^{++}$  nonet. The  $f_1(1510)$  was seen in  $K^-p \rightarrow \Lambda K^*\bar{K}\pi$  at 4 GeV/c [49], and at 11 GeV/c [50]. Evidence is also reported in  $\pi^-p$  at 8 GeV/c, based on the phase motion of the  $1^{++}$   $K^*\bar{K}$  wave [43]. A somewhat broader  $1^{++}$  signal is also observed in  $J/\psi(1S)$  radiative decay into  $\eta\pi^+\pi^-$  [51].

The absence of  $f_1(1420)$  in  $K^-p$  [50] argues against the  $f_1(1420)$  being the  $s\bar{s}$  member of the  $1^{++}$  nonet. However, the  $f_1(1420)$  was reported in  $K^-p$  but not in  $\pi^-p$  [52], while two experiments do not observe the  $f_1(1510)$  in  $K^-p$  [52,53]. The latter is also not seen in radiative  $J/\psi(1S)$  decay [8,9] and possibly [10], central collisions [47], or  $\gamma\gamma$  collisions [54], although, surprisingly for an  $s\bar{s}$  state, a signal is reported in  $4\pi$  decays [55]. These facts lead to the conclusion that  $f_1(1510)$  is not well established [56].

Assigning the  $f_1(1420)$  to the  $1^{++}$  nonet, one finds a nonet mixing angle of  $\sim 50^\circ$  [56]. However, arguments favoring the  $f_1(1420)$  being a hybrid  $q\bar{q}g$  meson, or a four-quark state, were put forward in Refs. [57] and [58], respectively, while Ref. [59] argued for a molecular state formed by the  $\pi$  orbiting in a  $P$ -wave around an  $S$ -wave  $K\bar{K}$  state.

Summarizing, there is convincing evidence for the  $f_1(1420)$  decaying into  $K^*\bar{K}$ , and for two pseudoscalars in the  $\eta(1440)$  region, the  $\eta(1405)$  and  $\eta(1475)$ , decaying into  $a_0(980)\pi$  and  $K^*\bar{K}$ , respectively. The  $f_1(1510)$  is not well established.

## References

1. P.H. Baillon *et al.*, Nuovo Cimento **50A**, 393 (1967).
2. D.L. Scharre *et al.*, Phys. Lett. **97B**, 329 (1980).
3. C. Edwards *et al.*, Phys. Rev. Lett. **49**, 259 (1982).
4. J.E. Augustin *et al.*, Phys. Rev. **D42**, 10 (1990).
5. J.Z. Bai *et al.*, Phys. Lett. **B594**, 47 (2004).
6. M.G. Rath *et al.*, Phys. Rev. **D40**, 693 (1989).
7. T. Adams *et al.*, Phys. Rev. Lett. **87**, 041801 (2001).

8. J.Z. Bai *et al.*, Phys. Rev. Lett. **65**, 2507 (1990).
9. J.E. Augustin and G. Cosme, Phys. Rev. **D46**, 1951 (1992).
10. M. Ablikim *et al.*, Phys. Rev. **D77**, 032005 (2008).
11. A. Bertin *et al.*, Phys. Lett. **B361**, 187 (1995).
12. A. Bertin *et al.*, Phys. Lett. **B400**, 226 (1997).
13. C. Cicalo *et al.*, Phys. Lett. **B462**, 453 (1999).
14. F. Nichitiu *et al.*, Phys. Lett. **B545**, 261 (2002).
15. T. Bolton *et al.*, Phys. Rev. Lett. **69**, 1328 (1992).
16. C. Amsler *et al.*, Phys. Lett. **B358**, 389 (1995).
17. S. Fukui *et al.*, Phys. Lett. **B267**, 293 (1991).
18. D. Alde *et al.*, Phys. Atom. Nucl. **60**, 386 (1997).
19. J.J. Manak *et al.*, Phys. Rev. **D62**, 012003 (2000).
20. B. Aubert *et al.*, Phys. Rev. Lett. **101**, 091801 (2008).
21. E. Klempt, Int. J. Mod. Phys. **A21**, 739 (2006).
22. E. Klempt and A. Zaitsev, Phys. Reports **454**, 1 (2007).
23. A.V. Anisovich *et al.*, Nucl. Phys. **A690**, 567 (2001).
24. A. Abele *et al.*, Phys. Rev. **D57**, 3860 (1998).
25. C. Amsler *et al.*, Eur. Phys. J. **C33**, 23 (2004).
26. F. Close *et al.*, Phys. Lett. **B397**, 333 (1997).
27. T. Barnes *et al.*, Phys. Rev. **D55**, 4157 (1997).
28. T. Gutsche *et al.*, Phys. Rev. **D79**, 014036 (2009).
29. M. Acciarri *et al.*, Phys. Lett. **B501**, 1 (2001).
30. F. Close *et al.*, Phys. Rev. **D55**, 5749 (1997).
31. D.M. Li *et al.*, Eur. Phys. J. **C28**, 335 (2003).
32. R. Ahohe *et al.*, Phys. Rev. **D71**, 072001 (2005).
33. P. Achard *et al.*, JHEP **0703**, 018 (2007).
34. G.S. Bali *et al.*, Phys. Lett. **B309**, 378 (1993).
35. L. Faddeev *et al.*, Phys. Rev. **D70**, 114033 (2004).
36. H.-Y. Cheng *et al.*, Phys. Rev. **D79**, 014024 (2009).
37. G. Li *et al.*, J. Phys. G: Nucl. Part. Phys. **35**, 055002 (2008).
38. T. Gutsche *et al.*, Phys. Rev. **D80**, 014014 (2009).
39. A. Masoni, C. Cicalo, and G.L. Usai, J. Phys. **G32**, R293 (2006).
40. C. Dionisi *et al.*, Nucl. Phys. **B169**, 1 (1980).
41. S.U. Chung *et al.*, Phys. Rev. Lett. **55**, 779 (1985).
42. D.F. Reeves *et al.*, Phys. Rev. **D34**, 1960 (1986).
43. A. Birman *et al.*, Phys. Rev. Lett. **61**, 1557 (1988).

- 44. H.J. Behrend *et al.*, Z. Phys. **C42**, 367 (1989).
- 45. A. Ando *et al.*, Phys. Rev. Lett. **57**, 1296 (1986).
- 46. T.A. Armstrong *et al.*, Phys. Lett. **B221**, 216 (1989).
- 47. D. Barberis *et al.*, Phys. Lett. **B413**, 225 (1997).
- 48. T.A. Armstrong *et al.*, Z. Phys. **C52**, 389 (1991).
- 49. P. Gavillet *et al.*, Z. Phys. **C16**, 119 (1982).
- 50. D. Aston *et al.*, Phys. Lett. **B201**, 573 (1988).
- 51. J.Z. Bai *et al.*, Phys. Lett. **B446**, 356 (1999).
- 52. S. Bitjukov *et al.*, Sov. J. Nucl. Phys. **39**, 738 (1984).
- 53. E. King *et al.*, Nucl. Phys. (Proc. Supp.) **B21**, 11 (1991).
- 54. H. Aihara *et al.*, Phys. Rev. **D38**, 1 (1988).
- 55. D.A. Bauer *et al.*, Phys. Rev. **D48**, 3976 (1993).
- 56. F.E. Close and A. Kirk, Z. Phys. **C76**, 469 (1997).
- 57. S. Ishida *et al.*, Prog. Theor. Phys. **82**, 119 (1989).
- 58. D.O. Caldwell, *Hadron 89 Conf.*, p. 127.
- 59. R.S. Longacre, Phys. Rev. **D42**, 874 (1990).